

CHANDRA X-RAY OBSERVATION OF THE ORION NEBULA CLUSTER. I. DETECTION, IDENTIFICATION, AND DETERMINATION OF X-RAY LUMINOSITIES

E. FLACCOMIO¹

Dipartimento di Scienze Fisiche ed Astronomiche, Università di Palermo, I-90123 Palermo, Italy;
ettoref@oapa.astropa.unipa.it

F. DAMIANI, G. MICELA, AND S. SCIORTINO

Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Palermo G. S. Vaiana,
Palazzo dei Normanni, I-90134 Palermo, Italy

AND

F. R. HARNDEN, JR., S. S. MURRAY, AND S. J. WOLK

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

Received 2002 March 4; accepted 2002 September 3

ABSTRACT

In this first of two companion papers on the Orion Nebula Cluster (ONC), we present our analysis of a 63 ks *Chandra* HRC-I observation that yielded 742 X-ray detections within the $30' \times 30'$ field of view. To facilitate our interpretation of the X-ray image, here we collect a multiwavelength catalog of nearly 2900 known objects in the region by combining 17 different catalogs from the recent literature. We define two reference groups: an *infrared sample*, containing all objects detected in the *K* band, and an *optical sample* comprising low-extinction, well-characterized ONC members. We show for both samples that field object contamination is generally low. Our X-ray sources are primarily low-mass ONC members. The detection rate for optical sample stars increases monotonically with stellar mass from zero at the brown dwarf limit to $\sim 100\%$ for the most massive stars but shows a pronounced dip between 2 and $10 M_{\odot}$. We determine L_X and L_X/L_{bol} for all stars in our optical sample and utilize this information in our companion paper to study correlations between X-ray activity and other stellar parameters.

Subject headings: open clusters and associations: individual (Orion Nebula Cluster) — X-rays: stars

On-line material: machine-readable tables

1. INTRODUCTION

The Orion region is one of the most frequently observed areas of the sky. It comprises several molecular clouds and stellar associations, among which the Orion Nebula cluster (ONC; also referred to as the Trapezium region) is of particular interest. With an age of about 1 Myr, more than 2000 young stellar objects with masses between 0.08 and $50 M_{\odot}$, an estimated central density of $\sim 2 \times 10^4$ stars parsec⁻³, and at a distance of ~ 470 pc, it is the largest and densest concentration of young pre-main-sequence (PMS) stars in our region of the Galaxy. Only part of the ONC members—those lying on the near side of the Orion molecular cloud from which the cluster is forming—are optically visible; more than half are so deeply embedded in the cloud that they are observable only at infrared (IR) or X-ray wavelengths, where the cloud becomes more transparent. For more complete descriptions of the ONC structure, dynamics, and stellar content, see Hillenbrand (1997), Hillenbrand & Hartmann (1998), Hillenbrand & Carpenter (2000), Carpenter (2000), Luhman et al. (2000), O'Dell (2001), and references therein. Among others, Hillenbrand and collaborators (Hillenbrand 1997; Hillenbrand et al. 1998; Hillenbrand & Carpenter 2000) have extensively studied both the optically visible and the embedded ONC stellar content.

Several studies (Stassun et al. 1999; Herbst et al. 2000; Herbst, Bailer-Jones, & Mundt 2001; Rebull 2001) have measured rotational periods via photometric modulation of stellar spots. Spectacular *Hubble Space Telescope* (*HST*) images (see, e.g., Bally, O'Dell, & McCaughrean 2000) have allowed the direct observation of many circumstellar disks and *proplyds* (photoionization structures due to the evaporation of disks) that seem to surround a large fraction of the PMS stars in the vicinity of the central bright star θ^1 Ori C.

The ONC was first detected in X-rays by the *Uhuru* satellite (Giacconi et al. 1972). Subsequent imaging observations performed with the *Einstein* and *ROSAT* observatories (Ku & Chanan 1979; Gagné, Caillault, & Stauffer 1995) indicated that (1) ONC members are powerful X-ray emitters, with typical L_X/L_{bol} close to the saturation value, 10^{-3} , observed for fast rotators on the main sequence; (2) no relation seems to hold between X-ray activity and stellar rotation; and (3) stars with circumstellar accretion disks may have lower levels of X-ray activity. These works were, however, hindered by the lack, at that time, of complete optical information on the ONC population and by the low sensitivity and spatial resolution of the X-ray instrumentation, especially important to resolve the dense central region and to unambiguously identify X-ray sources with optical counterparts. Both of these limitations have been recently lifted: *Chandra* observations (Schulz et al. 2001; Garmire et al. 2000) have revealed about 1000 X-ray point sources, for the most part associated with low-mass stars down to the substellar mass limit.

¹ Current address: Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Palermo G. S. Vaiana, Palazzo dei Normanni, I-90134 Palermo, Italy.

In this work we describe the analysis of a *Chandra* High Resolution Camera (HRC; Murray et al. 2000) observation of the ONC and correlate our data with relevant data from the literature. In a companion work (Flaccomio et al. 2003, hereafter Paper II), we investigate statistical correlations between X-ray activity and other stellar characteristics (such as rotational period, mass, age, and disk accretion indicators) and search for insights into the physical mechanisms that drive activity in PMS stars. An investigation of the X-ray variability characteristics of ONC members based on these data is in preparation.

The structure of this paper is as follows: We first introduce our *Chandra* observation and discuss its reduction and X-ray source detection (§ 2). We then describe the preparation of an extensive catalog of known X-ray/optical/IR/radio objects in the region of our HRC observation, collecting from the literature useful observable properties and defining two reference object samples (§ 3). In the same section we also report the results of our observation in the BN/KL region. In § 4 we concentrate on the optical/IR properties of our X-ray sources, and in § 5 and a supporting appendix, we determine the X-ray activity indicators L_X and L_X/L_{bol} , along with their uncertainties. Our findings are summarized in § 6, setting the stage for Paper II.

2. OBSERVATION AND DATA REDUCTION

The HRC on board the *Chandra X-Ray Observatory* (Weisskopf et al. 2002) observed the ONC for 63.2 ks on 2000 February 4. The pointing (R.A. = $5^{\text{h}}35^{\text{m}}17^{\text{s}}$, decl. = $-5^{\circ}23'16''$ [J2000.0]) was chosen to place the Trapezium region and the bright O star θ^1 Ori C in the center of the field of view (FOV). A good fraction of the ONC region was included in the $30' \times 30'$ HRC FOV.

Our *Chandra* observations offer these significant improvements over previous *Einstein* and *ROSAT* observations: unprecedented spatial resolution ($\sim 0''.5$ in the field center), important in such a crowded field (see § 3.2); sensitivity ~ 20 times deeper than previous X-ray data (see § 5); and continuous temporal coverage for variability studies.

2.1. Data Filtering

Our HRC analysis begins by filtering the event list to lower background levels² and increase the source signal-to-noise ratio (SNR). We considered filtering out particular time intervals and/or ranges of pulse height amplitude (PHA) dominated by instrumental and spacecraft environment effects. We examined light curves of the entire observation but found that our particular data set did *not* suffer from variable solar activity contamination commonly reported for *Chandra* data. Hence, no time filtering was applied. On the other hand, PHA distributions of all events do show a sharp instrumental peak for PHA values ≤ 35 , and as in our previous HRC analyses (Harnden et al. 2001), we excluded this peak by eliminating events with PHA < 35 . Given the PHA response of the HRC to X-rays and stellar spectra (see § 5.1), these counts are almost exclusively background: total events were reduced by about 15% (from ~ 2.1 to $\sim 1.8 \times 10^6$), while source counts declined only $\sim 1\%$.

² Comprising $\sim 80\%$ of the total count rate, background dominates the data.




Fig. 1.—HRC-I image of the ONC processed with CSMOOTH. Linear gray scale indicates photon arrival rate vs. position on the detector. In this low-resolution presentation, black indicates scale saturation due to intense pointlike sources (or unresolved groups of sources). Note that the PSF increases significantly toward image edges.

Figure 1 shows the filtered HRC image, smoothed for presentation purposes with the CSMOOTH task in the *Chandra* Interactive Analysis of Observations (CIAO) package.³

2.2. Source Detection

We have analyzed the filtered event list with the Palermo wavelet transform detection code, PWDETECT (Damiani et al. 1997; F. Damiani et al. 2003, in preparation).⁴ In synthesis, PWDETECT analyzes an image at a variety of spatial scales, allowing detection of both pointlike and moderately extended sources. The most important parameter required by the code is the detection threshold which determines, along with the image background level, the number of expected spurious detections. The relationship between detection threshold and expected number of spurious sources was determined by running the PWDETECT code on 400 simulated background images with roughly the same number of photons as in the background of our observation ($\sim 1.46 \times 10^6$, once the source contribution is removed).

We choose to accept detections with $\text{SNR} > 4.74$, corresponding to 10 expected spurious detections throughout the FOV. We accept 10 spurious sources, instead of the more customary one, for three reasons: (1) 10 is still a small fraction of the total number of detected sources; (2) by lowering the threshold from 5.25σ (corresponding to one spurious source) to 4.74, we gain ~ 100 good detections (after a “hand filtering” of the detection list; see below), of which only nine are expected to be spurious; and (3) as shown in § 3.2, *Chandra*’s superb spatial resolution ensures that the ~ 10

³ Available at <http://asc.harvard.edu/~ciao>.

⁴ See also http://www.astropa.unipa.it/progetti_ricerca/PWDetect.

TABLE 1
X-RAY SOURCES DETECTED IN THE ORION NEBULA CLUSTER

N_X	R.A.	Decl.	$\delta_{R.A., decl.}$	SNR	Counts	δ_{counts}	Exposure Time
(1)	(J2000.0)	(J2000.0)	(arcsec)	(5)	(6)	(7)	(ks)
	(2)	(3)	(4)				(8)
1.....	5 34 14.46	-5 28 16.62	3.38	36.27	1277.75	67.97	49.06
2.....	5 34 18.23	-5 33 28.53	6.91	6.65	167.54	41.78	47.64
3.....	5 34 20.70	-5 30 46.81	5.90	9.21	259.90	46.23	49.44
4.....	5 34 20.74	-5 23 25.85	8.03	4.76	111.66	33.69	51.22
5.....	5 34 24.96	-5 22 06.32	3.17	42.89	1095.36	46.16	52.24
6.....	5 34 27.23	-5 24 22.23	2.02	63.91	1866.58	64.05	52.90
7.....	5 34 27.69	-5 31 55.39	3.17	42.05	1536.32	72.59	50.23
8.....	5 34 28.53	-5 24 58.95	2.20	50.53	1225.93	49.62	53.21
9.....	5 34 29.09	-5 14 32.51	4.46	15.73	434.32	225.67	49.69
10.....	5 34 29.30	-5 23 56.09	2.30	45.86	1024.30	45.09	53.52

NOTES.—Units of right ascension are hours, minutes, and seconds, and units of declination are degrees, arcminutes, and arcseconds. Table 1 is published in its entirety in the electronic edition of the *Astrophysical Journal*. A portion is shown here for guidance regarding its form and content.

spurious sources are extremely unlikely to be identified with other wavelength counterparts.

Visual inspection of the initial lists of ~ 830 detections produced by PWDETECT showed that although the vast majority of detections appear real and most of the obvious sources in the image are detected, two “glitches” of the detection algorithm produce undesired effects. First, about 90 sources are detected in the vicinity of very bright stars, notably on the wings of the point-spread function (PSF) of the central source (θ^1 Ori C). These are most likely spurious detections due to large intensity fluctuations on the wings of bright sources. In such regions the code is unable to estimate correctly the highly spatially variable background level. Second, in a few cases (~ 20) a pair of sources clearly visible and distinguishable by eye in the X-ray image is detected as both a point source corresponding to one of the two and an extended source encompassing the pair. We dealt with the first problem (to be addressed in a future version of PWDETECT) by visually examining the list of detections and excluding the spurious sources in question. Specifically, we deleted all sources lying closer than $6''8$ to θ^1 Ori C, thereby also losing any real sources occurring in that area. The second problem occurs because the extended detection has a higher SNR than the pointlike detection. For such cases, we eliminated this problem by running a special version of PWDETECT that limits its search to pointlike sources.

PWDETECT estimates positions and count rates of detected sources in the assumption that the PSF is Gaussian. Since the actual HRC PSF is not Gaussian, we investigated the effect of this simplifying assumption on our estimates. We computed the wavelet transform of calibration PSFs (in the *Chandra* Calibration Database)⁵ at several detector positions and for a representative energy of ~ 1.5 keV. In order to better match the actual shape of bright sources in our HRC image, these calibration PSFs were first convolved with a bi-dimensional Gaussian with $\sigma = 0''3$, possibly reflecting the effect of imperfect aspect reconstruction affecting the real source PSF. Corrections with respect to the Gaussian PSF approximation were parameterized as a function of off-axis angle and, for count rates, ranged from

16% at the field center to 0% at $\gtrsim 10'$ off-axis. Having verified that this result agrees well with the (background subtracted) extracted counts of bright isolated sources in the limit of large extraction radii, we applied the correction to our source count rates. Positional errors (up to $0''9$ for off-axis angles of $\sim 14'$) are instead always much smaller with respect to random uncertainties and were therefore neglected.

The final lists of detections, comprising 742 X-ray sources, is presented in Table 1, where we give the sky position and its statistical uncertainty, detection SNR, number of source counts and uncertainty, and effective exposure time at the source positions. This latter quantity, describing the spatially varying sensitivity of the HRC plus *Chandra* mirror system, is derived from an exposure map calculated with CIAO for an incident energy of 2.0 keV, i.e., the approximate temperature of our sources (see § 5.1). This choice of temperature is not critical for our purposes, however, because the normalized effective area at any given point on the detector depends only marginally on energy: for $0.5 < kT < 3.5$, the values of effective exposure times at any given location on the detector vary at most at the $\sim 4\%$ level.

In addition to measuring count rates for detected X-ray sources, we also used PWDETECT to compute upper limits to the photon flux of all nondetected objects in the catalog described in the next section. This calculation was performed consistently using the same SNR threshold as for detections and applying the correction, described above, due to the non-Gaussian PSF.

3. THE GENERAL CATALOG OF KNOWN OBJECTS

We assembled an extensive catalog of known X-ray/optical/IR and radio objects that fell within the HRC FOV. In addition to our list of HRC sources and the *Chandra* source lists of Garmire et al. (2000) and Schulz et al. (2001), we considered 14 catalogs from recent publications, producing a database of nearly 2900 distinct objects reported in at least one of the studies considered. A full list of references is given in the first column of Table 2 along with a concise classification of the work (col. [2]) and the referenced table number(s) from the original work (col. [3]).

⁵ See <http://asc.harvard.edu/caldb>.

TABLE 2
LIST OF CATALOGS IN THE ONC AREA

Reference (1)	Type of Study (2)	Table (3)	Number ^a (4)	R_{95} ^b (5)	N_R ^c (6)	N_{HRC} ^d (7)	Data ^e (8)
This work.....	X-ray (HRC-I)	1	742	2.2 ^f	553	...	HRC-I count rate
Schulz et al. 2001.....	X-ray (ACIS-S)	1	111	0.4	63	92	ACIS-S count rate
Garmire et al. 2000.....	X-ray (ACIS-I)	1	971 ^g	1.4	559	579	...
Bally et al. 2000.....	Optical (<i>HST</i>)	1, 2	48	0.5	17	14	...
Hillenbrand 1997.....	Optical phot./spec.	1, 3	1516	0.4	1055	627	$V^1 J^2 A_V^2 T_{\text{eff}} L_{\text{bol}}$
Hillenbrand et al. 1998 ^h	IR photometry	1	1516	627	$J^2 H^3 K^2 L^2 \text{EW}_{\text{Ca II}} \Delta_{I-K}$
Hillenbrand & Carpenter 2000 ⁱ	IR photometry	1	776	0.4	242	282	$H^1 K^1$
Lucas & Roche 2000.....	IR photometry	1	557	0.7	102	116	$I^1 J^1 H^2$
Lucas et al. 2001.....	IR spectroscopy	1, 3	23	2	A_V^1 mass
Lada et al. 2000.....	IR phot. protostars	1	78	0.5	11	26	L^1
2MASS.....	IR photometry	...	1596	0.6	728	416	$J^4 H^5 K^4$
Carpenter et al. 2001.....	IR phot./variability	4, 7	555	0.5	379	241	$J^3 H^4 K^3 P_{\text{rot}}^4$
Stassun et al. 1999.....	P_{rot}	1	124	0.5	115	84	P_{rot}^3
Herbst et al. 2000 ^j	P_{rot}	2	132	109	P_{rot}^2
Herbst et al. 2001 ^k	P_{rot}	...	296	168	P_{rot}^1
Jones & Walker 1988.....	Proper motion	3	1034	519	P_{memb}
Felli et al. 1993.....	Radio	1, 5	49	0.9	18	25	...
M. Reid 2002, private communication.....	Radio	...	100	0.4	32	43	...

^a Number of sources in the HRC field of view.

^b The 95% quantiles, in arcseconds, of the object distances from the Jones & Walker 1988 counterparts.

^c Number of objects used to derive R_{95} , i.e., identified with Jones & Walker 1988.

^d Number of objects identified with an HRC X-ray source.

^e Data collected from each catalog; superscripts give precedence for acceptance in our catalog.

^f In the central $5' \times 5'$ of the FOV, where the *Chandra* PSF is narrower, $R_{95} \sim 0''.65$, based on 148 objects.

^g The original lists contains 973 sources. We deleted two that had repeated coordinates.

^h Same catalog as in Hillenbrand 1997.

ⁱ Covers only the central $5' \times 5'$ of the HRC FOV.

^j Subset of the catalog of Jones & Walker 1988.

^k Data provided by W. Herbst. The original list contained 403 stars; we used data only for those in common with Jones & Walker 1988.

3.1. Catalog Cross-Identifications

Cross-identifications were based on positional coincidence. First of all, we registered the coordinates of each catalog to a standard reference list (that of Jones & Walker 1988) using the mean relative displacement of well-identified objects. In the case of the Garmire et al. (2000) list of X-ray sources, we also had to apply a rotation of $0^\circ:21$ around the HRC field center in order to bring the coordinates into agreement with those of the other lists. We then performed the final identifications using the following tolerances: (1) For our list of HRC X-ray sources, we used the positional uncertainties computed by PWDETECT as a function of source statistics and off-axis angle. These range from $0''.036$ for the bright central source (θ^1 Ori C) to $\sim 8''$ for weak sources close to the detector border. (2) For the *Chandra* sources of Garmire et al. (2000), uncertainties were assumed to follow the same off-axis trend as measured in our data. We performed a fourth-order polynomial fit of HRC position errors versus off-axis angle and then conservatively used twice these values for the Garmire et al. (2000) source off-axis angles (to allow for additional counting statistics effects). (3) For all other catalogs, we used ~ 2.5 times the mean dispersion of the registered position offsets with respect to the reference list (Jones & Walker 1988). In all cases, this identification radius was $\lesssim 1''.0$.

We then examined the resulting list of identifications by eye and modified some identifications according to more subjective criteria. For example, nine X-ray sources detected at large off-axis ($\geq 8'$), and therefore with relatively large PSFs, were associated with one or two optical objects that

fell outside the error circle. In these cases these optical objects lay well within the PSF and thus potentially contributed the detected photons. In some of these cases the source centroid lay between two objects unresolved (by PWDETECT), so neither was within the formal error circle. In all such cases we added the missing identifications manually.

The end result is a table (not shown) with 15 columns, one for each catalog,⁶ and 2887 rows, each row representing the cross-identifications of a single object. Table 2 gives, for each catalog, the number of objects in the HRC FOV (col. [4]). Column (5) gives a measure of the relative uncertainty in the catalog coordinates, R_{95} , representing the 95% quantile of the object's offsets from the stars of the reference catalog (Jones & Walker 1988). Column (6) reports the number of objects used in the latter calculation. With typical values of $R_{95} \sim 0''.5$, the number of uncertain or ambiguous identifications is indeed small. The number of objects in each catalog identified with an HRC X-ray source is given in column (7). In the next section we discuss in detail the important issue of the reliability and uniqueness of these HRC source identifications.

3.2. X-Ray Source Identification

The outstanding sharpness of the *Chandra* PSF, especially in the field center, makes it possible for the first time to identify detected X-ray sources as easily as can be done in

⁶ Although Table 2 lists 17 catalogs, two of these refer to one of the other lists for object identification and are therefore not counted in the number of cross-identified catalogs.




FIG. 2.—Cumulative distributions of offsets between X-ray and optical positions, for three optical catalogs: Jones & Walker (1988), Hillenbrand (1997), and Hillenbrand & Carpenter (2000). For the first two, we show separate distributions for the inner $5' \times 5'$ and for the entire FOV. Since the third catalog covers only the central region, a single distribution is shown.

the optical and IR passbands, even in an extremely crowded field like the ONC. This fact largely eliminates the identification uncertainties that were unavoidable with all previous X-ray observatories and makes assignment of physical parameters for the X-ray emitters much easier than before. Figure 2 shows the cumulative distribution of the offsets of our HRC sources with respect to three of our component catalogs. Offsets near the field center are particularly small and depend systematically on the comparison catalog, the positions of Hillenbrand & Carpenter (2000) being most similar to *Chandra*'s. It is therefore quite likely that optical/IR uncertainties are an important (if not predominant) contribution to the field-center offset distribution. Although identification radii are small, the surface density of objects in our catalog is extremely high, especially in the central $\sim 5' \times 5'$ Trapezium region where both the actual density of ONC members and the sensitivity of the optical/IR catalogs are highest. Fortunately, the field center is also where the HRC PSF is sharpest.

In order to verify that the probability of spurious identification is small, we calculated the expected number, $N_{\text{ch.id}}$, of chance correlations in a given region of our image by estimating the object surface density at the positions of our sources (ρ_{obj}), multiplying these densities by the coincidence region areas (A_{id}),⁷ and then summing over all sources, i.e.,

$$N_{\text{ch.id}} = \sum_i A_{\text{id}}^i \rho_{\text{obj}}^i, \quad (1)$$

⁷ For this calculation, we took a quadrature sum of X-ray position uncertainty and a generous optical uncertainty of $0''.75$ as the identification radius.

where the sum is extended over all sources in the region of interest. To estimate the surface density ρ_{obj} , we characterized the FOV in two different ways: as square tiles of $1'$ on a side and as a series of concentric annuli centered on the X-ray field, with the radius of each annulus 0.5 greater than that of the previous one. We then counted the number of cataloged sources in each of these regions and divided by their area. The two methods gave equivalent results. The probability of chance identification is highest in the field center, $\sim 5\%$ for the inner $1'$ radius; for the entire field, the chance probability is $\sim 2\%$. If anything, these probabilities could even be smaller, since actual identification radii for most optical catalogs were smaller than the conservative $0''.75$ used here.

We can compare these estimates with the number of spurious X-ray sources expected on the basis of the SNR threshold chosen for detection (see § 2.2). With 10 such spurious detections expected in the full FOV, none (i.e., $\lesssim 0.2$) will be associated by chance with a known object. This means we need not worry about spurious sources when studying the properties of previously cataloged objects. When exploring the nature of unidentified sources (see Paper II), however, we must remember that ~ 10 detections will be spurious.

We can also estimate the number of times an X-ray source identified with its true counterpart will be identified by chance with a second object. Assuming that the object positions are uncorrelated, we multiply the fraction of chance identifications (2%) by the total number of sources (742) to get an expected $\lesssim 15$ double chance identifications in the entire field.⁸ Since we actually find 50 multiple identifications, the source positions probably are correlated.

3.3. Physical Properties of Cataloged Stars

For each object in our merged catalog, we collected relevant data such as X-ray count rates, optical and IR photometry, rotational periods, etc. In cases of redundant information, only one of the available values was chosen: the last column of Table 2 reports the information used from each catalog, and in cases of redundant information, a superscript on a quantity indicates the precedence rank with which values were adopted for our merged list (e.g., H^5 was used only if the other four catalogs with H values had no entry for a given object). Table 3 gives the number of objects in our catalog for which we retrieved several measured or estimated quantities, as well as the same number referred to HRC sources identified with a single optical/IR object. The most complete data for objects are the near-IR H and K magnitudes, since the majority of the ONC population is embedded in the molecular cloud and is only detectable at IR wavelengths where the extinction ($A_K \sim 0.1A_V$) is greatly reduced from that in the optical.

A first look at our stars via IR photometry is shown in Figure 3 as a K versus $H-K$ diagram for the 763 stars in the inner $5' \times 5'$ of the FOV. The same diagram for the entire HRC FOV would be similar, but, with 2431 stars, it would

⁸ With chance double identifications scaling as the square of the radius, and for X-ray errors of $\gtrsim 2''$ typical of *XMM-Newton* and the best prior studies (Gagné et al. 1995), the expected fraction of double identifications in the central $1'$ would be $\gtrsim 45\%$ (for *Chandra*-level sensitivity and assuming uncorrelated object positions). Even neglecting source confusion problems, no other past or present X-ray telescope is suitable for ONC study.

TABLE 3
NUMBERS OF OBJECTS WITH EACH OPTICAL/IR DATA ITEM

Sample	Spectral Type	A_V	L_{bol}	T_{eff}	Mass	Age	P_{rot}	Member ^a
Whole sample	1001	924	907	908	900	873	451	1028
HRC detections	460	437	437	437	434	427	242	492
Sample	Δ_{I-K}	EW _{Ca II}	V	I	J	H	K	L
Whole sample	785	872	1176	1625	2339	2532	2466	132
HRC detections	389	385	501	596	604	642	641	49

^a Indicates stars with measured proper motion.

be overcrowded. X-ray sources are depicted with filled symbols, and optically well-characterized stars are shown with larger, open circles. Also shown are the main sequence and a 1 Myr isochrone derived by transforming the evolutionary tracks of Siess, Dufour, & Forestini (2000, hereafter SDF) from the theoretical (L_{bol} vs. T_{eff}) to the observational plane using the transformations of Bessell (1991) and Houdashelt, Bell, & Sweigart (2000). Reddening vectors for 1 Myr stars of three different masses are indicated by dashed lines. We cannot derive precise information on the nature of our stars from this diagram alone: H - and K -band magnitudes are indeed expected to be influenced by the presence of disks around many of our stars. We expect that stars with disks will be displaced with respect to those without disks toward the upper right of Figure 3. For 813 stars in common with our sample, Hillenbrand et al. (1998) derive a mean⁹ I_c-K

⁹ The corresponding median value is 0.26 mag.




FIG. 3.—Near-IR color-magnitude diagrams for the central $5' \times 5'$ of the HRC FOV. Large and small symbols indicate stars belonging and not belonging to the optical sample (see § 3.4). Filled and empty symbols indicate objects unambiguously detected and not detected, respectively, in the HRC data. Also shown are the main sequence and the 1 Myr isochrone according to SDF evolutionary tracks.

color excess of 0.36 mag, with a standard deviation of 0.72 mag. Given that disk emission is expected to be minimum at I -band wavelengths, these values can be taken as a rough measure of the average K -band excess and as an upper limit to the $H-K$ color excess. We can conclude that the positions of stars in Figure 3, although influenced by excesses, are approximately representative of their photospheres. We then observe that our near-IR catalog contains stars down to very low masses ($M \ll 0.1 M_{\odot}$) and high extinction ($A_V > 50$) and that stars are detected in our HRC data down to $M \gtrsim 0.1 M_{\odot}$ and to quite high extinction ($A_V \sim 40$).

One of our main goals, to be pursued further in Paper II, is to study the activity of young stars as a function of their mass and evolutionary stage. The most relevant data in this context are those given in the papers by Hillenbrand (1997) and Hillenbrand et al. (1998). The largest part of our spectral types, extinction estimates (A_V), and T_{eff} and L_{bol} measurements are taken from Hillenbrand (1997), and we refer the reader to that work for details. The same author also derives stellar masses and ages using the D'Antona & Mazzitelli (1994) evolutionary tracks. Since new improved model calculations have been published, we decided to estimate masses and ages anew from the published temperatures and luminosities. We considered two sets of calculations: those by D'Antona & Mazzitelli (1997, hereafter DM97;¹⁰ $Z = 0.02$, $Y = 0.28$, $X_d = 2 \times 10^{-5}$) and those of SDF ($Z = 0.02$, $Y = 0.277$, $X = 0.703$ and no overshooting). Recent estimates of dynamical masses for seven single stars and two binary systems by Simon, Dutrey, & Guilloteau (2000) seem to favor the SDF over the DM97 tracks. We also observe that for ONC members the SDF tracks produce a smaller age spread than do the DM97 calculations, possibly better reflecting the real star formation history. Given uncertainties in the models, it is premature to rely exclusively on a particular set of tracks. We primarily have adopted the SDF tracks and in Paper II will note whether and how our results are influenced by this choice over that of the DM97 calculations.

Masses and visual absorptions for 19 presumed brown dwarfs ($0.008 \leq M/M_{\odot} \leq 0.08$) and three low-mass stars studied with IR spectroscopy by Lucas et al. (2001) were used when available. For four of these objects we could also derive masses using the optical data of Hillenbrand (1997) and the SDF tracks: three are brown dwarfs ($M_{\text{IR}} = 0.073$, 0.074, and 0.071 M_{\odot}) according to the IR data and very

¹⁰ The authors issued an update in 1998 for $M < 0.9 M_{\odot}$, available on line at <http://www.mporzio.astro.it/~dantona/prems.html>.

low mass stars ($M_{\text{opt}} = 0.11, 0.11, \text{ and } 0.19 M_{\odot}$) according to the optical estimate; the fourth is a very low mass star with $M_{\text{IR}} = 0.3 M_{\odot}$ and $M_{\text{opt}} = 0.21 M_{\odot}$.

3.4. Definition of Stellar Samples

About a third (907) of the cataloged objects can be placed on the H-R diagram and belong to the well-characterized population studied by Hillenbrand (1997) and Hillenbrand et al. (1998). We were able to estimate masses and ages for 877 of these stars that lie within the SDF tracks. Masses for nine massive stars ($M/M_{\odot} > 6.0$) were taken from Hillenbrand (1997). For four stars in the SDF tracks (see § 3.3) and for 18 other very low mass stars and brown dwarfs, we adopt values by Lucas et al. (2001). For the most part, the remaining cataloged objects (~ 2000) are heavily obscured stars detected only in the near-IR and for which we therefore have limited information.

Hillenbrand (1997) argues that her sample, which comprises most of our well-characterized sample, is representative of the whole ONC population and differs only in that its location on the near side of the molecular cloud gives it low optical extinction. This is important for our study of X-ray activity (Paper II) because it ensures that our results will not be strongly biased. As expected in a flux-limited sample, however, the brighter and more massive stars studied by Hillenbrand (1997) are actually seen to larger values of visual extinction than their lower mass counterparts. In order to minimize selection effects for our studies, we limited part of our analysis to stars with $A_V < 3.0$.

For the following analysis we define two groups of stars, mainly on the basis of the amount of available information:

1. Our *optical sample* is comprised of stars in the HRC FOV for which we have a mass estimate, whose values of A_V are less than 3.0, and which are either confirmed proper motion members or have unknown proper motion (see § 3.4.1). For the 696 stars of this optical sample, Table 4 lists sky position, mass, age, rotational period, Ca II line equivalent width, HRC *basal* count rate (see § 5), X-ray luminosity, and L_X/L_{bol} (see §§ 4 and 5). The analysis of X-ray activity that we present in Paper II is based on these data.

2. The *IR sample* is comprised of 2476 stars with measured *K*-band magnitudes. The IR sample includes most of

the optical sample, with 680 of 696 stars in common. This sample very likely contains a large fraction of all the ONC members of stellar and brown dwarf mass, with the possible exception of a few deeply reddened stars located in small regions where the cloud absorbing column is greatest (see Paper II). The central region, which also coincides with a thicker part of the molecular cloud, has been surveyed in depth by Hillenbrand & Carpenter (2000) and Lucas & Roche (2000). The survey of Hillenbrand & Carpenter (2000) is believed better than 90% complete to $K \sim 17.5$, corresponding to a mass of $\sim 0.02 M_{\odot}$ (for an age of 1 Myr).

Figure 4 shows an H-R diagram for the stars in our optical sample. We also show the positions of other confirmed proper motion members ($P_{\text{memb}} > 50\%$) that are not part of the optical sample. For both groups we indicate stars (see legend in figure) that are unambiguously identified with our X-ray sources. SDF tracks and isochrones used to estimate stellar masses and ages are also shown for reference.

In the following two sections we estimate the degree to which our two main stellar samples are contaminated by field objects not physically or spatially related to the ONC.

3.4.1. Contamination of the Optical Sample

In discussing the optical sample and its H-R diagram, we will make use of the membership information provided by Jones & Walker (1988). We consider three stellar subsamples that differ in confidence of their physical association with the ONC. The first, group 0, comprises all stars studied astrometrically by Jones & Walker (1988), regardless of membership probability as derived in that study. The next sample, group 1, comprises all stars with proper motion membership probabilities greater than 50%. Third, we consider the optical sample defined in the previous section, which excludes only those stars explicitly suspected of being nonmembers ($P_{\text{memb}} < 50\%$).

We can estimate the number of field objects, N_{field} , that contaminates a (sub)sample of stars studied astrometrically (Jones & Walker 1988) by interpreting proper motion probabilities literally: $N_{\text{field}} = \sum (1.0 - P_i)$, where P_i is the probability that the i th object belongs to the ONC and the sum is extended over the whole (sub)sample. Table 5 displays our results, with the first two rows referring to stars in

TABLE 4
CATALOG OF THE ONC OPTICAL SAMPLE

N	R.A.	Decl.	Mass	$\log(\text{age})$	P_{rot}	EW_{CaII}	Count Rate ^a	$\log(L_X)$	$\log(L_X/L_{\text{bol}})$
(1)	(J2000.0)	(J2000.0)	(M_{\odot})	(yr)	(days)	(7)	(ks^{-1})	(ergs s^{-1})	(10)
1.....	5 34 12.81	-5 28 48.28	0.72	6.46	...	1.60	<4.02	<30.24	<-3.14
2.....	5 34 14.39	-5 28 16.77	1.74	6.87	...	2.00	27.39	31.21	-3.12
3.....	5 34 17.15	-5 29 04.45	0.23	4.71	...	0.00	<3.28	<30.57	<-3.08
4.....	5 34 17.92	-5 33 33.45	0.43	6.04	3.19	1.80	4.59	30.23	-3.23
5.....	5 34 19.39	-5 27 12.04	1.86	6.79	...	4.00	<2.83	<30.04	<-4.35
6.....	5 34 19.47	-5 30 19.99	1.19	6.20	...	-2.50	<3.62	<30.48	<-3.42
7.....	5 34 20.64	-5 32 35.24	0.19	6.29	...	1.80	<3.75	<30.14	<-2.70
8.....	5 34 20.72	-5 23 29.18	0.28	6.55	...	0.00	1.74	29.81	-2.96
9.....	5 34 20.92	-5 24 48.53	0.25	6.52	...	0.70	<2.56	<29.97	<-2.78
10.....	5 34 22.32	-5 22 27.01	0.12	5.07	<2.40	<29.95	<-3.08

NOTES.—Units of right ascension are hours, minutes, and seconds, and units of declination are degrees, arcminutes, and arcseconds. Table 4 is published in its entirety in the electronic edition of the *Astrophysical Journal*. A portion is shown here for guidance regarding its form and content.

^a Basal HRC count rate (see text).




FIG. 4.—H-R diagrams for stars belonging to the optical sample and for other ONC members that we could place in the diagram; see legend in figure for symbol meanings. We indicate with filled circles and small plus signs the stars in these two groups that were detected in the HRC image. Dashed and solid lines, respectively, represent SDF evolutionary tracks for masses from 0.1 to 7.0 M_{\odot} and isochrones for ages between 10^5 and 10^8 yr (note labels beside lines).

the entire HRC FOV and the inner $5' \times 5'$ region. The remainder of the table reports the full-field analysis as a function of stellar mass and age. The table gives total numbers of objects and field contamination fractions estimated for the stars in group 0 (P_{memb} known; cols. [2] and [3]) and

group 1 ($P_{\text{memb}} \geq 50\%$; cols. [4] and [5]). The last two columns refer to the optical sample, which contains both stars with $P_{\text{memb}} \geq 50\%$ and stars with no proper motion information. For the former stars we estimate the fraction of field objects using the same procedure as for groups 0 and 1; for the latter, we assume that this fraction is the same as for group 0 (i.e., we use the values from col. [3]).

Regarding sample contaminations, we therefore draw the following conclusions: (1) Field contamination for the whole sample is a factor of ~ 2 lower ($\sim 5\%$) in the central region with respect to the full FOV ($\sim 10\%$), most likely because of the high member surface density and the optically opaque molecular cloud that effectively obscures background stars. (2) Selection of stars with $P_{\text{memb}} \geq 50\%$ (i.e., group 1) reduces the contamination to 2%–2.5%, regardless of the area considered. (3) Contamination depends on position in the H-R diagram, being highest for masses between 1 and 10 M_{\odot} and for the oldest stars [$\log(\text{age}) > 6.5$] in group 0; this is likely the result of both the distribution of field stars in the observational H-R diagram and of the dependence on mass and age of the member sky density (the 3–10 M_{\odot} stars, for example, are the most uniformly distributed among the mass ranges considered and, given the centrally peaked member distribution, are found in regions of higher field star relative density). (4) Contamination in the optical sample is probably not much larger than in group 1, our estimate being 3.5% for the full FOV and 2.4% for the central region. We therefore take the optical sample as our primary basis for the analysis of Paper II.

3.4.2. Contamination of the IR Sample

For stars detected only in the K band, we cannot perform the above estimates because of lack of proper motions or any other individual membership data. Nonetheless, it is particularly important to have at least a statistical understanding of the contamination, since we expect deep K -band observations to begin to see through the dense molecular cloud.

TABLE 5
FIELD STAR CONTAMINATION FOR VARIOUS SAMPLES

Sample (1)	$N(0)^a$ (2)	Field Percentage (0) (3)	$N(1)^b$ (4)	Field Percentage (1) (5)	$N(\text{Opt.})^c$ (6)	Field Percentage (Opt.) ^d (7)
Whole FOV	1034	10.9	938	2.4	696	3.5
Inner $5' \times 5'$	241	4.6	235	2.2	138	2.4
Mass and Age Breakdown						
$M/M_{\odot} = 0.10\text{--}0.16$	43	4.5	42	2.2	61	2.9
$M/M_{\odot} = 0.16\text{--}0.25$	135	5.6	129	1.6	160	2.4
$M/M_{\odot} = 0.25\text{--}0.50$	302	5.1	291	1.8	277	2.2
$M/M_{\odot} = 0.50\text{--}1.00$	115	9.8	106	2.5	89	2.8
$M/M_{\odot} = 1.00\text{--}2.00$	75	26.8	56	3.5	40	5.1
$M/M_{\odot} = 2.00\text{--}3.00$	52	15.1	46	4.3	36	4.7
$M/M_{\odot} = 3.00\text{--}10.00$	20	22.6	16	5.8	12	6.7
$M/M_{\odot} = 10.00\text{--}50.00$	6	5.3	6	5.3	6	5.3
$\log(\text{age}) = 4.50\text{--}5.50$	61	5.3	59	2.1	56	2.6
$\log(\text{age}) = 5.50\text{--}6.00$	136	8.4	126	1.9	117	2.6
$\log(\text{age}) = 6.00\text{--}6.50$	366	5.6	352	2.3	330	2.5
$\log(\text{age}) = 6.50\text{--}7.50$	167	18.7	139	2.7	162	6.4

^a Number of stars with known proper motion (Jones & Walker 1988).
^b Number of stars with $P_{\text{memb}} \geq 50\%$.
^c Number of stars in the optical sample.
^d Extrapolated fraction of field contamination (see text).

We estimate the expected contamination by field objects (Galactic and extragalactic) with the following procedure. Briefly, we first determine a reasonable approximation to field objects' surface density as a function of unabsorbed K magnitude. Since the majority of field objects will be in the background with respect to the ONC, their density at a given magnitude will be lowered by absorption in the molecular cloud. A smaller, unabsorbed fraction will be in the foreground. We will assume that the shape of the K distribution for the two populations is the same as that of the total field population. Second, in order to know the total number of contaminating nonmembers, we must estimate the ratio of the number of foreground to background stars. Third, we use the ^{13}CO measurements published by Goldsmith, Bergin, & Lis (1997) to estimate the total molecular cloud absorption in K (A_K) as a function of sky position. Note that because these data are available only in the field center the contamination can be estimated only in the central region. Fourth, we convolve the background star K distribution with the A_K distribution over the area of interest, and fifth, we add the unmodified foreground star distribution to obtain the expected density of the field stars as a function of K .

In the first step of the calculation outlined above, the field star density as a function of K was derived from more than 13,600 stars in the 2MASS survey¹¹ within 1° of the Trapezium: although this distribution is affected by the presence of the cluster and of the absorbing cloud, the vast majority of stars from which it is derived come from the field and are located off the cloud, so that the final result will not be significantly affected. It is important to realize what the limiting magnitude of our K -band luminosity function is. Although the nominal 2MASS survey completeness limit in K is 14.3, in many cases it goes fainter than this, and for the 1° radius region considered here, the K -band luminosity function rises to $K \sim 15$ and turns over for fainter K magnitudes. We will therefore take $K = 15$ as our limit to the knowledge of the K distribution. Considering that more than 90% of the area of interest has a ^{13}CO column density corresponding to $A_K > 1.0$, we can confidently estimate background contamination down to $K \sim 16$.

In the second step above, where we estimate the fraction of foreground stars in the direction of the ONC and in a given solid angle, $f_{\text{ONC}} = N_{\text{ONC}}(d < 470 \text{ pc})/N_{\text{ONC}}(\text{tot})$, we use the results obtained by Carpenter (2000) through the Wainscoat et al. (1992) Galactic model. Although the author does not report a result for the ONC, he quotes a ratio of foreground to total field stars for Perseus, $f_{\text{Per}} = N_{\text{Per}}(d < 320 \text{ pc})/N_{\text{Per}}(\text{tot})$, that has a Galactic latitude ($b \sim -20^\circ$) similar to that of the ONC and a distance of 320 pc (vs. 470 pc). We assume that the object space density as a function of distance, $\rho(d)$, is the same in the two directions: with this assumption, the fraction of foreground stars in the ONC can be expressed as

$$f_{\text{ONC}} = f_{\text{Per}} \frac{\langle \rho(470) \rangle}{\langle \rho(320) \rangle} \left(\frac{470}{320} \right)^3 \lesssim f_{\text{Per}} \left(\frac{470}{320} \right)^3, \quad (2)$$

where $\langle \rho(d) \rangle$ is the mean stellar density within a distance d and the last inequality is justified by the larger height of the ONC above the Galactic plane. According to Carpenter

(2000), f_{Per} ranges from 0.05 to 0.08 in the J , H , and K bands. We have (conservatively) taken the upper end of this range and scaled it by cube of the distances (eq. [2]), therefore deriving $f_{\text{ONC}} \lesssim 0.25$. We will conservatively assume in the following that $f_{\text{ONC}} = 0.25$; this could result in an overestimation of field contamination.

The fifth step of our calculation gives us the expected surface density of field stars in the central $5' \times 5'$ region as a function of magnitude. For example, we find a value of 0.16 stars arcmin⁻² for $K < 13$ and 0.44 stars arcmin⁻² for $K < 15$. These numbers can be compared with the observed mean densities in the same region at the same two limiting magnitudes: ~ 21 and ~ 28 stars arcmin⁻² for $K < 13$ and $K < 15$, respectively. Even for $K < 16$, the expected contamination from field stars is about 2%.

Having estimated the field star surface densities, we now compare our determinations with others, performed with similar data and in a similar fashion. Consider the densities of unabsorbed field stars. Carpenter, Hillenbrand, & Skrutskie (2001), in their work on IR variability, derive an *off-cloud* density of 0.66 stars arcmin⁻² for $K < 14.8$. Carpenter (2000) performs a similar calculation of the unabsorbed field star densities for the ONC and other clouds. From the author's Figure 3, we can see that the mean field star density for $K < 14.3$ at the Galactic latitude of the ONC (i.e., -19°) is about 0.5. The densities that we derive are 0.81 for $K < 14.8$ and 0.64 for $K < 14.3$, both within 20%–30% of the previous derivations. Because our goal is to show that contamination is not significant, a more conservative approach is to adopt our slightly larger derivation for the stellar surface density. Comparing our expectation for the density of observed field stars with the results of Hillenbrand & Carpenter (2000), we find that their densities are about 50%–60% lower than ours, in keeping with our conservative approach.

We conclude, therefore, that contamination in our IR sample is negligible in the center of the FOV down to $K \sim 16$. Note that this conclusion might not hold for the whole area under study, because of the lower average member densities and thinner background molecular cloud. The IR photometry in the outer regions is not as deep as in the center, however, so that even there contamination is possibly not significant.

3.5. The BN/KL Region

The BN/KL region, about $1'$ northwest of the Trapezium, in the densest part of Orion Nebula cloud, contains a cluster of massive embedded stars mostly observed in the IR. The X-ray emission of the IR sources in the region has been already investigated by Garmire et al. (2000) using *Chandra* Advanced CCD Imaging Spectrometer-I (ACIS-I) data. We estimate, using PIMMS,¹² that our HRC-I data is 1.5 to 4.0 times less sensitive, depending on the source kT and N_{H} (see § 5.1), with the worst case corresponding to high absorptions ($N_{\text{H}} \sim 10^{23}$). With reference to the sky region depicted in Figure 6 of Garmire et al. (2000), our master catalog contains 61 distinct objects in the ~ 0.6 arcmin² area, 22 of which are detected by the HRC, versus 27 by ACIS (Table 2 in Garmire et al. 2000). Out of the five ACIS sources missed by the HRC, three are classified as IR

¹¹ See <http://www.ipac.caltech.edu/2mass>.

¹² Portable, Interactive, Multi-Mission Simulator, Version 3.0, available on line at <http://asc.harvard.edu/toolkit/pimms.jsp>.

sources and two as X-ray-only sources by Garmire et al. (2000). We do not detect the BN object itself, but our upper limit (<12 photons in 63 HRC-I ks) is consistent with the low measured ACIS-I count rate (18 photons in 48 ACIS-I ks). The IR source “n” (our source 272) likely varied between the time of the two observations: the HRC detects 49 photons while ACIS detects 61, which, assuming $N_H = 10^{23}$ (Garmire et al. 2000) and $kT > 1$ keV, indicates that the HRC luminosity is ≥ 3 times greater than that measured by ACIS. The IR source 17 in Table 2 of Garmire et al. (2000), our source 281 in Table 1, was, with similar assumptions (but $N_H = 10^{22.5}$), 6 times brighter in the HRC data. Two objects detected exclusively in X-rays also appear to have varied: number 16 (HRC source 284), ~ 3 times brighter in the HRC, and number 23, undetected by the HRC and whose upper limit indicates a variation of at least a factor of 3.

4. OPTICAL/IR PROPERTIES OF THE HRC SOURCES

In this section we discuss the K magnitude and estimated mass composition of the counterparts of our X-ray source sample. In doing so, we often distinguish the two samples of optically characterized members and near-IR objects. In Paper II, following our study of correlations of X-ray activity with stellar parameters, we speculate on the possible nature of X-ray sources not identified with optical or IR counterparts.

The quantity most widely available for our cataloged objects is K -band magnitude. In the upper panel of Figure 5 we show the K -band magnitude distribution for all our cataloged objects together with that of the subsample of stars comprising our optical sample. Also shown are distribu-




FIG. 5.—Upper panel shows K -band distributions for the IR sample (i.e., all objects within the HRC FOV with measured K magnitudes; black), X-ray-detected and unambiguously identified objects (light gray), the well-characterized optical sample (hatched; see text), and optical sample objects detected in X rays (dark gray). Lower panel displays HRC detection fractions for the IR and optical samples (light and dark gray, respectively) as a function of K magnitude.

TABLE 6
DETECTION FRACTIONS VERSUS K MAGNITUDE

Sample	Sky region	$K \leq 11$	$K \leq 14$
IR sample.....	Full FOV	0.57	0.32
	Central $5' \times 5'$	0.63	0.41
Optical sample	Full FOV	0.69	0.50
	Central $5' \times 5'$	0.81	0.71
Optical sample (max) ^a	Central $5' \times 5'$	0.90	0.77

^a Including ambiguous identifications.

tions referring to objects in these two samples that are detected in our HRC data and unambiguously identified with an IR or optical counterpart. The lower panel of the figure shows detection fractions of the two reference samples as a function of K . It can be seen that the fraction of objects detected in X-rays is greater than 50% down to $K \sim 11$ but then decreases for fainter K magnitudes, going to zero at $K \sim 14$, which roughly corresponds to the brown dwarf limit (see Fig. 3) for 1 Myr-old unabsorbed stars. Similar results, although with slightly larger detection fractions, were obtained by Garmire et al. (2000) with their ACIS-I data. Optical sample stars follow a similar trend, although with detection fractions somewhat larger. A quantitative breakdown of detection fractions for several stellar samples and two limiting K magnitudes is given in Table 6. Note the increased detection fractions in the center of the FOV.

Figure 6 (*upper panel*) shows the distribution of stellar masses for our optical sample and the subset of X-ray detected stars. As in the previous figure, the lower panel shows the detection fraction: we see that the detection fraction is 100% for the most massive stars, drops abruptly for $\log(M/M_\odot) = 0.3-1.0$ ($M/M_\odot \sim 2.0-10.0$),




FIG. 6.—Upper panel: Mass distributions of objects in the HRC FOV and belonging to our optical sample. Shadings same as in Fig. 5. Lower panel: Detection fraction vs. mass; error bars computed from Poisson statistics.

rises again to about 90%, and then decreases fairly smoothly to zero in the brown dwarf regime. A similar trend (not shown) is observed in the field center, although the detection fractions are 10%–20% higher between ~ 0.1 and $\sim 2.0 M_{\odot}$.

5. DETERMINATION OF L_X AND L_X/L_{bol} FOR HRC SOURCES

Here we estimate two activity indicators, X-ray luminosity (L_X) and its ratio to bolometric luminosity (L_X/L_{bol}), and estimate their associated uncertainties. L_X represents total coronal power output, while L_X/L_{bol} is related to the fraction of total stellar energy that goes

into heating the corona. In our search for physical mechanisms that may drive activity in young PMS stars, Paper II studies the relationship of these indicators with various stellar parameters.

Because the intensity of many of the HRC sources varied during our observation, we did not use mean count rates to estimate luminosities. Instead, we have defined a *basal* count rate to remove the effects of short-term ($\lesssim 63$ ks) variability. Our basal rate algorithm searches for that count rate compatible with the largest possible portion of the light curve. Figure 7 shows five light curves of highly variable sources and their basal rates, and Table 4 lists our estimates for the basal count rates, L_X , and the ratio between L_X and L_{bol} for the stars in our optical sample.




Fig. 7.—Light curves for five strongly variable HRC sources (number indicated at upper left of each panel) vs. time elapsed since observation start. Bin size varies from source to source, and left and right ordinate scales show count rates and counts per bin, respectively. Solid horizontal lines indicate derived basal rates (see text); dashed lines show zero levels. Source data were extracted from circular regions around each position, with subtracted background estimated from larger concentric annuli.

5.1. *Chandra* HRC Count-Rate-to-Flux Conversion Factors

In order to convert basal count rates to intrinsic X-ray luminosity (assuming a distance of 470 pc), we have derived conversion factors (CFs) that yield unabsorbed source flux from observed count rate. We have assumed both that emitted radiation can be described as thermal, optically thin emission from an isothermal plasma at temperature kT and that the absorbing interstellar material along the line of sight can be characterized by a hydrogen column density N_H . We find that the X-ray flux of stars in our optical sample can be adequately described by an isothermal ($kT = 2.16$), optically thin plasma, absorbed by a gas column density proportional to optical extinction: $N_H = 2 \times 10^{21} A_V$. This conclusion is based upon a spectral analysis of archival ONC X-ray data from ACIS (Townsend et al. 2000), as discussed in the Appendix. Because the *Chandra* HRC lacks spectral resolution, N_H and kT values cannot be deduced from our main data set alone.

In order to study the sensitivity of the conversion factor to the assumed plasma temperature and hydrogen column density, we have explored the two-dimensional (kT , N_H) parameter space in the range $kT < 8.0$ and $20 < \log(N_H) < 23$, which most likely includes the conditions of the vast majority of our sources. All our HRC CFs are computed using PIMMS, a Raymond-Smith spectral model, and a spectral band between 0.1 and 4.0 keV.¹³ Figure 8 shows a contour plot of the ratio of the conversion factor at a given kT and N_H within the specified ranges to the conversion factor for typical ONC values of $kT = 2.16$ and $N_H = 3 \times 10^{21}$. The plot shows that the dependence of the conversion factor on temperature and N_H is quite weak

¹³ This band was chosen to ease comparisons with previous *Einstein* and *ROSAT* results. A different choice would lead to different conversion factors depending on source temperature: for the full *Chandra* band (0.2–8.0) the fractional difference in CFs compared to our choice would be between +7.5% ($kT = 1.3$) and –20% ($kT = 3.5$), with the most likely value $\sim -4.5\%$ ($kT = 2.16$).




Fig. 8.—Contour plot of the ratio between the HRC conversion factor computed for $kT = 2.16$ and $N_H = 3 \times 10^{21}$ (2.9×10^{-11} ergs cm^{-2} photon $^{-1}$; large triangle) and the same conversion factor at different values of kT and N_H . Right vertical axis gives the standard conversion between N_H and A_V . Region between the two dashed vertical lines depicts likely range of ONC source temperatures.

for $kT > 1.0$ and values of N_H around the assumed value. The weak dependence of the CF on temperature justifies (at least for purposes of computing the CFs) the assumption of a single temperature instead of the full emission measure distribution. Provided the majority of emitting plasma is at temperatures larger than ~ 1.0 keV (likely for ONC PMS stars) and the estimates of N_H are reasonable, our approach will give useful results.

The next step, for each of our sources, is to estimate a temperature representative of the emitting plasma and a value of N_H quantifying the amount of interstellar gas between the source and the observer. If the range of such representative temperatures is such that $kT > 1.0$ keV (see § A2), we may safely use a single temperature (~ 2 keV) to compute the CFs for all sources. Although this kT is somewhat higher than 1 keV, a value often used in past X-ray studies of PMS stars, the HRC conversion factors we thus derive differ, for typical N_H , by at most 0.1 dex. N_H , on the other hand, must be evaluated individually since it certainly spans a large range of values. We can estimate N_H independently from X-ray information by using the *standard* linear correlation with the optical absorption: $N_H = 2.0 \times 10^{21} A_V$ (see Ryter 1996). Adopting such a linear relation between optical absorption (due to dust grains) and gas column density (responsible for X-ray absorption) assumes that the gas-to-dust ratio is the same for all of our sources and equal to the value inferred for field stars. This assumption might not hold for the ONC members for two reasons: (1) most of the absorption is due to the molecular cloud in which the cluster is embedded rather than to interstellar matter between us and the cloud, and these two mediums might have different gas-to-dust ratios; and (2) the circumstellar environment of these PMS stars, many of which are actively accreting material from disks, is complex and might in principle contribute to an anomalous extinction law. An independent check on the validity and limits of the above assumption (see the Appendix) allows us to proceed in using this N_H - A_V relationship to derive N_H from the A_V values of Hillenbrand (1997) for many of our ONC members. This is particularly applicable for stars in our optical sample that enter into the analysis of correlations between activity and stellar parameters (Paper II).

To summarize this discussion, as well as that of the Appendix, the conversion between count rates and energy fluxes or luminosities is a crucial step, usually requiring some spectral information. We have satisfied this need with archival ACIS data and conclude that our conversion factors are good to better than $\sim 30\%$ (0.1 dex), albeit with a 1σ scatter of ~ 0.2 dex. This conclusion does not appear to depend on stellar mass or accretion rate.

5.2. Uncertainties on L_X and L_X/L_{bol}

Uncertainties in our inferred X-ray luminosities are predominately due to the conversion from counts to flux (except for the weakest of our sources, where Poisson statistics are also important). In § 5.1 we found that the statistical uncertainties on the CFs will be of the order of 0.2 dex (1σ). Count rate measurements have a median fractional Poisson uncertainty of $\sim 17\%$ (+0.07, –0.08 dex), small compared to the total L_X error, but for 5% (or 35%) of sources, this component is $\geq 50\%$ (or 30%), comparable to the conversion factor uncertainty.

Distance uncertainties will also influence luminosity values. At 470 pc, the ONC's angular extent ($\sim 30'$) corresponds to ~ 4 pc, and the range of modern distance determinations introduces an additional systematic uncertainty of ~ 30 pc, together yielding $\Delta L_X \approx 13\%$. Since such largely systematic uncertainties will not qualitatively influence our results for the dependence of coronal activity on stellar parameters, we neglect these.

The L_X/L_{bol} ratio will be affected by uncertainties in both L_X and L_{bol} . For stars in our optical sample, we have adopted the L_{bol} values of Hillenbrand (1997) with typical errors of $\lesssim 0.2$ dex (stars with strong circumstellar accretion could have slightly larger, underestimation errors). A quadrature sum of the uncertainties of ~ 0.2 dex for $\log(L_X)$ and 0.2 dex for $\log(L_{\text{bol}})$ yields a total $\log(L_X/L_{\text{bol}})$ uncertainty of ~ 0.3 dex.

6. SUMMARY

Our PWDETECT wavelet algorithm finds 742 X-ray sources in the $30' \times 30'$ FOV of our 63 ks *Chandra* HRC-I observation of the ONC. As a basis for our study of ONC X-ray activity (see Paper II), we have combined 17 different catalogs from the recent literature to assemble a catalog of all available information for nearly 2900 objects, the large majority of which are ONC members. We then calculated mass and age for a significant subset using the evolutionary tracks of Siess, Dufour, & Forestini (2000) and defined two reference stellar samples: the *optical sample*, comprising ~ 700 well-characterized members with low extinction ($A_V \leq 3.0$), and the *IR sample*, largely including the optical

sample and comprising ~ 2500 stars observed in the *K* band. We have shown that field object contamination is quite limited in both of these samples.

In order to characterize the population of X-ray detected objects, we have presented distributions of *K* magnitudes and masses and have noted, in agreement with the results of Gagné et al. (1995) and Garmire et al. (2000), a clear dependence of the percentage of detections on stellar mass, from $\sim 0\%$ at the brown dwarf limit to $\sim 100\%$ for $\sim 2.0 M_\odot$ stars. For $2\text{--}10 M_\odot$ stars, the detection fraction drops sharply, followed by a marked increase (to 100%) for the six highest mass ONC stars.

We concluded here by estimating activity indicators (L_X and L_X/L_{bol}) for stars in our optical sample; Paper II studies correlations between this X-ray activity and other stellar parameters. Our X-ray luminosities were computed from basal count rates, assuming single-temperature spectra for all sources and proportionality between X-ray absorption and optical extinction, with these assumptions verified by our analysis of medium-resolution archival ACIS X-ray spectra for a subset of our sources.

The authors would like to thank M. Reid for providing his catalog of radio sources in the ONC area prior to publication. This work was partially supported at the CfA by NASA contracts NAS8-38248 and NAS8-39073 and by NASA grant NAS5-4967. F. D., E. F., G. M., and S. S. wish to acknowledge support from the Italian Space Agency (ASI) and MURST. E. F. would like to thank the CfA for its hospitality during his Fellow visits.

APPENDIX A

THE ACIS DATA

On 1999 October 12–13, ACIS (Townsend et al. 2000) observed the central region of the ONC for a total exposure time of ~ 47 ks (observation ID 18, sequence 200016). These data were discussed by Garmire et al. (2000) and are available in the *Chandra* archive.

Here we utilize ACIS medium-resolution spectra to demonstrate that the fluxes of our HRC sources can be adequately described by an isothermal ($kT = 2.16$), optically thin plasma, absorbed by a gas column density proportional to optical extinction: $N_{\text{H}} = 2 \times 10^{21} A_V$. This verifies the assumptions of § 5 for a range of temperatures ($kT > 0.5$ keV) and for adherence to the relationship between X-ray absorbing column and optical extinction.

Our spectral analysis characterized source spectra with two hardness ratios, HR_1 and HR_2 , and compared the measured HR pairs with model predictions for a grid of kT and N_{H} values. This approach is justified by the fact that large statistical uncertainties for the majority of our sample¹⁴ mask any (smaller) systematic inaccuracies introduced by our simplifying assumptions.

A1. DATA PROCESSING

In order to improve upon the standard energy calibration, we applied the CORRECT_CTI procedure (Townsend et al. 2000) to the level 1 event file. This procedure (partially) corrects for the spatial nonuniformity of the ACIS spectral response that stems from degradation of the charge transfer inefficiency (CTI) during the early days of the *Chandra* mission. With this correction, spectral resolution is $\sim 100\text{--}200$ eV, depending both on the incidence position with respect to the readout node and on the photon energy. Following this correction we applied standard grade and status flag filtering to produce a “clean” level 2 event file.

We then extracted background-subtracted source counts in three spectral bands: $0.5\text{--}1.7$ keV (*L*), $1.7\text{--}2.8$ keV (*M*), and $2.8\text{--}8.0$ keV (*H*), for 678 HRC sources that fell in the ACIS FOV. Source spectra were extracted from circles centered on the HRC source positions, with a radius 3 times the HRC source size as measured by PWDETECT (see § 2.2). Background was estimated from concentric annuli of inner and outer radii of 4 and 5 times the HRC source size. To avoid contamination

¹⁴ For the brightest sources a more detailed spectral analysis can be performed.

effects, we discarded 36 sources because of the proximity (distance less than 6 times the size) of another HRC source, and 150 additional sources were excluded due to net counts ≤ 0 in one or more of the three spectral bands.

For the remaining 492 sources, we calculated two hardness ratios defined as $HR_1 = (M - L)/(M + L)$ and $HR_2 = (H - M)/(H + M)$. Uncertainties on the hardness ratios were derived assuming that the number of source and background photons in each spectral band followed a Poisson distribution with mean equal to the measured value.

A2. INTERPRETATION

We computed hardness ratio predictions for a grid of isothermal Raymond-Smith models with a range of temperatures and hydrogen column densities, using PIMMS.¹⁵ Figure 9 shows the results of our PIMMS calculations in a grid of HR_1 versus HR_2 ; also shown are positions of high-SNR HRC sources (see legend). By interpolation in this plane, we can derive kT^{HR} and N_{H}^{HR} , where the superscript refers to the method used to estimate these quantities. From these two spectral parameters we can derive the desired HRC conversion factors, CF^{HR} (see Fig. 8).

Formal 1σ errors on kT^{HR} , N_{H}^{HR} , and CF^{HR} were derived (along with uncertainties on the hardness ratios) by Monte Carlo methods with the assumption that extracted source (and background) counts were Poisson distributed with mean equal to the measured value. In order to compute mean values and errors, we require that the source has at least a 90% chance of lying inside the HR grid. For this reason a few points that lie inside the grid but close to its boundaries are excluded from the subsequent analysis, along with those that lie outside. These latter points may be explained either by statistical fluctuation in the hardness ratio or by the fact that their spectra cannot be modeled as simply as we have assumed (e.g., they have a significant second temperature thermal component; this could explain the points on the left-hand side of Fig. 9).

The majority of our sources have low SNR, in both the HRC and the ACIS data sets. This affects our hardness ratio analysis: the 1σ error bars are in general quite large. Examination of Figure 9 shows that (especially at very low and very high values) kT^{HR} and N_{H}^{HR} are quite strong functions of the two ratios, making it difficult to determine the two parameters. We therefore decided to include in the following conversion factor analysis only sources with errors on HR_1 and HR_2 less than

¹⁵ Since PIMMS ignores the finite spectral resolution of ACIS, we chose spectral bands to minimize the potential effect of significant spectral features appearing in the selected passband boundaries. To confirm that we have indeed avoided this problem, we computed focal plane spectra for nonabsorbed sources for each grid temperature, convolved these spectra with Gaussians of varying FWHM, and then compared the hardness ratios obtained from these smoothed spectra to those obtained from unsmoothed spectra. For an FWHM of 300 eV, the difference between the two HRs is less than 0.023 in all cases, negligible compared to typical uncertainties in our measured HR values.




FIG. 9.—Grid of predicted ACIS hardness ratios for absorbed single-temperature Raymond-Smith models; kT ranges from 0.27 to 8.62 keV, with $\log(N_{\text{H}})$ running from 20 to 23.3. Filled circles denote optical sample stars that have hardness ratios and errors within the grid and whose inferred values of kT , N_{H} , and HRC conversion factor were used for our analysis (see text). Open circles have HR values (or errors) near the grid edge, preventing their use. Plus signs denote positions (and error bars) of HRC sources with computed errors of less than 0.1 for HR_1 and HR_2 .




FIG. 10.—Ratio of HRC count rate to flux conversion factors vs. mass, computed from optical extinction and $kT = 2.16$ keV as described in the text. Only stars with high SNR are shown, with vertical error bars propagated from Poisson statistics. Horizontal segments with diamonds in their center represent median values of the points in the corresponding mass ranges (as employed in Paper II).

0.1, i.e., the subsample depicted in Figure 9. This choice is dictated by our interest in trends of median CF^{HR} with mass, accretion, etc. By definition, the median is not weighted and does not take into account variable errors, and including a larger number of low-SNR points would not necessarily improve the statistical significance of the median.

The median kT^{HR} for our high-SNR group is 2.2 keV, with 90% of the sources in the 1.6–3.5 keV range. A fraction of this scatter can be explained by uncertainties on the hardness ratios: with the mean temperature as reference (i.e., ~ 2.4 keV), the formal χ^2 is only 2.8. We examined trends in kT with stellar mass and age, finding no significant evidence for a dependence in the range $0.2\text{--}10 M_{\odot}$. Neither could we find any significant difference in the kT distributions of “high” and “low” accretion stars as defined from the Ca II equivalent width (see Paper II). Given the range of temperatures found and the discussion concerning Figure 8 (see § 5.1), we can be assured that an error on the assumed kT does not influence the HRC conversion factor by more than $\sim 10\%$. Consequently, the most important aspect of our analysis becomes the study of the relation between N_H^{HR} and A_V .

The mean ratio between A_V and N_H^{HR} is 0.54×10^{-21} , not very different from the nominal value 0.5×10^{-21} . However, there are several objects that lie far from this mean relation, so that the quartiles of the ratio distribution are 0.21×10^{-21} and 0.77×10^{-21} . It is unclear whether this large spread of values is due to a real departure from the N_H versus A_V mean relation (i.e., to an anomalous gas-to-dust ratio) or to other factors such as an oversimplified spectral model.

To conclude, we examine the most important relation for our purpose, i.e., the relation between conversion factors derived from hardness ratios, CF^{HR} , and those derived assuming a constant temperature ($kT = 2.16$) and N_H proportional to A_V (CF_{A_V}). The median for the logarithm of the CF^{HR}/CF_{A_V} ratio is -0.024 , and the 1σ quantiles are -0.252 and 0.130 . Figure 10 shows this ratio as a function of stellar mass for our high-SNR sample, as well as median values in the mass ranges defined in Paper II and depicted in the figure as horizontal segments. No significant trend in the median is apparent. We also considered separately samples of “high-accretion” and “low-accretion” stars, finding no difference in the conversion factor ratio; sample sizes for determining this latter comparison, however, are very small: nine “high-accretion” stars versus 14 with “low accretion” (see Paper II).

REFERENCES

- Bally, J., O’Dell, C. R., & McCaughrean, M. J. 2000, *AJ*, 119, 2919
 Bessell, M. S. 1991, *AJ*, 101, 662
 Carpenter, J. M. 2000, *AJ*, 120, 3139
 Carpenter, J. M., Hillenbrand, L. A., & Skrutskie, M. F. 2001, *AJ*, 121, 3160
 Damiani, F., Maggio, A., Micela, G., & Sciortino, S. 1997, *ApJ*, 483, 350
 D’Antona, F., & Mazzitelli, I. 1994, *ApJS*, 90, 467
 ———. 1997, *Mem. Soc. Astron. Italiana*, 68, 807 (DM97)
 Felli, M., Taylor, G. B., Catarzi, M., Churchwell, E., & Kurtz, S. 1993, *A&AS*, 101, 127
 Flaccomio, E., Damiani, F., Micela, G., Sciortino, S., Harnden, F. R., Murray, S. S., & Wolk, S. J. 2003, *ApJ*, 582, 398 (Paper II)
 Gagné, M., Caillault, J., & Stauffer, J. R. 1995, *ApJ*, 445, 280
 Garmire, G., Feigelson, E. D., Broos, P., Hillenbrand, L. A., Pravdo, S. H., Townsley, L., & Tsuboi, Y. 2000, *AJ*, 120, 1426
 Giacconi, R., Murray, S., Gursky, H., Kellogg, E., Schreier, E., & Tananbaum, H. 1972, *ApJ*, 178, 281
 Goldsmith, P. F., Bergin, E. A., & Lis, D. C. 1997, *ApJ*, 491, 615
 Harnden, F. R., et al. 2001, *ApJ*, 547, L141
 Herbst, W., Bailer-Jones, C. A. L., & Mundt, R. 2001, *ApJ*, 554, L197
 Herbst, W., Rhode, K. L., Hillenbrand, L. A., & Curran, G. 2000, *AJ*, 119, 261
 Hillenbrand, L. A. 1997, *AJ*, 113, 1733
 Hillenbrand, L. A., & Carpenter, J. M. 2000, *ApJ*, 540, 236
 Hillenbrand, L. A., & Hartmann, L. W. 1998, *ApJ*, 492, 540
 Hillenbrand, L. A., Strom, S. E., Calvet, N., Merrill, K. M., Gatley, I., Makidon, R. B., Meyer, M. R., & Skrutskie, M. F. 1998, *AJ*, 116, 1816
 Houdashelt, M. L., Bell, R. A., & Sweigart, A. V. 2000, *AJ*, 119, 1448
 Jones, B. F., & Walker, M. F. 1988, *AJ*, 95, 1755

- Ku, W. H.-M., & Chanan, G. A. 1979, *ApJ*, 234, L59
Lada, C. J., Muench, A. A., Haisch, K. E., Lada, E. A., Alves, J. F., Tollestrup, E. V., & Willner, S. P. 2000, *AJ*, 120, 3162
Lucas, P. W., & Roche, P. F. 2000, *MNRAS*, 314, 858
Lucas, P. W., Roche, P. F., Allard, F., & Hauschildt, P. H. 2001, *MNRAS*, 326, 695
Luhman, K. L., Rieke, G. H., Young, E. T., Cotera, A. S., Chen, H., Rieke, M. J., Schneider, G., & Thompson, R. I. 2000, *ApJ*, 540, 1016
Murray, S. S., et al. 2000, *Proc. SPIE*, 4012, 68
O'Dell, C. R. 2001, *PASP*, 113, 29
Rebull, L. M. 2001, *AJ*, 121, 1676
Ryter, C. E. 1996, *Ap&SS*, 236, 285
Schulz, N. S., Canizares, C., Huenemoerder, D., Kastner, J. H., Taylor, S. C., & Bergstrom, E. J. 2001, *ApJ*, 549, 441
Siess, L., Dufour, E., & Forestini, M. 2000, *A&A*, 358, 593 (SDF)
Simon, M., Dutrey, A., & Guilloteau, S. 2000, *ApJ*, 545, 1034
Stassun, K. G., Mathieu, R. D., Mazeh, T., & Vrba, F. J. 1999, *AJ*, 117, 2941
Townsend, L. K., Broos, P. S., Garmire, G. P., & Nousek, J. A. 2000, *ApJ*, 534, L139
Wainscoat, R. J., Cohen, M., Volk, K., Walker, H. J., & Schwartz, D. E. 1992, *ApJS*, 83, 111
Weisskopf, M. C., Brinkman, B., Canizares, C., Garmire, G., Murray, S. S., & Van Speybroeck, L. P. 2002, *PASP*, 114, 1